Location: Saltwater NP Survey method: Fixed walk around the site, duration 1.5-2 hours N° of species recorded: 129 Date this list was prepared: 26 April 2018 N° of surveys conducted (up to the preparation date): 84 Comment: No surveys are done in January because the site becomes crowded with campers



Alan Stuart

81 Queens Rd New Lambton NSW 2305 Australia Mobile: +61 409 978171 Email: <u>alanstuart400@gmail.com</u> Web: <u>www.thinkingaboutbirds.com</u>

## Species recorded, and their recording rates

| Black Swan                    | 4.8%  | Australian Pelican           | 19.1% |
|-------------------------------|-------|------------------------------|-------|
| Pacific Black Duck            | 1.2%  | Nankeen Night-Heron          | 25.0% |
| Chestnut Teal                 | 2.4%  | Striated Heron               | 9.5%  |
| Australian Brush-turkey       | 79.8% | Great Egret                  | 3.6%  |
| Brown Quail                   | 1.2%  | White-faced Heron            | 46.4% |
| White-headed Pigeon           | 27.4% | Little Egret                 | 6.0%  |
| Brown Cuckoo-Dove             | 4.8%  | Eastern Reef Egret           | 2.4%  |
| Wonga Pigeon                  | 42.9% | Australian White Ibis        | 1.2%  |
| Crested Pigeon                | 8.3%  | Australasian Gannet          | 14.3% |
| Peaceful Dove                 | 1.2%  | Little Pied Cormorant        | 9.5%  |
| Bar-shouldered Dove           | 21.4% | Great Cormorant              | 26.2% |
| Brown-capped Emerald-Dove     | 2.4%  | Little Black Cormorant       | 13.1% |
| Wompoo Fruit-Dove             | 9.5%  | Pied Cormorant               | 32.1% |
| Rose-crowned Fruit-Dove       | 1.2%  | Australasian Darter          | 1.2%  |
| Topknot Pigeon                | 10.7% | Osprey                       | 21.4% |
| Eastern Koel                  | 20.2% | Little Eagle                 | 1.2%  |
| Channel-billed Cuckoo         | 2.4%  | Grey Goshawk                 | 1.2%  |
| Shining Bronze-Cuckoo         | 10.7% | White-bellied Sea-Eagle      | 21.4% |
| Fan-tailed Cuckoo             | 17.9% | Whistling Kite               | 39.3% |
| Brush Cuckoo                  | 1.2%  | Brahminy Kite                | 10.7% |
| Tawny Frogmouth               | 1.2%  | Rainbow Bee-eater            | 3.6%  |
| White-throated Needletail     | 7.1%  | Dollarbird                   | 13.1% |
| Fork-tailed Swift             | 1.2%  | Azure Kingfisher             | 6.0%  |
| Beach Stone-curlew            | 1.2%  | Sacred Kingfisher            | 9.5%  |
| Australian Pied Oystercatcher | 48.8% | Laughing Kookaburra          | 91.7% |
| Sooty Oystercatcher           | 8.3%  | Nankeen Kestrel              | 1.2%  |
| Red-capped Plover             | 16.7% | Australian Hobby             | 1.2%  |
| Double-banded Plover          | 1.2%  | Yellow-tailed Black-Cockatoo | 1.2%  |
| Black-fronted Dotterel        | 6.0%  | Galah                        | 8.3%  |
| Masked Lapwing                | 13.1% | Australian King-Parrot       | 2.4%  |
| Bar-tailed Godwit             | 2.4%  | Eastern Rosella              | 1.2%  |
| Common Sandpiper              | 2.4%  | Rainbow Lorikeet             | 13.1% |
| Silver Gull                   | 50.0% | Scaly-breasted Lorikeet      | 41.7% |
| Little Tern                   | 10.7% | Noisy Pitta                  | 3.6%  |
| Australian Gull-billed Tern   | 6.0%  | Green Catbird                | 26.2% |
| Caspian Tern                  | 3.6%  | Regent Bowerbird             | 57.1% |
| Common Tern                   | 11.9% | Satin Bowerbird              | 25.0% |
| Crested Tern                  | 42.9% | White-throated Treecreeper   | 22.6% |
| Wedge-tailed Shearwater       | 3.6%  | Variegated Fairy-wren        | 29.8% |
|                               |       |                              |       |

| Superb Fairy-wren           | 65.5% | Olive-backed Oriole        | 3.6%  |
|-----------------------------|-------|----------------------------|-------|
| Red-backed Fairy-wren       | 1.2%  | Pied Currawong             | 28.6% |
| Scarlet Honeyeater          | 47.6% | Australian Magpie          | 83.3% |
| Noisy Friarbird             | 15.5% | Pied Butcherbird           | 51.2% |
| Brown Honeyeater            | 2.4%  | Grey Butcherbird           | 6.0%  |
| Crescent Honeyeater         | 1.2%  | White-breasted Woodswallow | 2.4%  |
| White-cheeked Honeyeater    | 45.2% | Spangled Drongo            | 36.9% |
| Eastern Spinebill           | 45.2% | Willie Wagtail             | 73.8% |
| Lewin's Honeyeater          | 96.4% | Rufous Fantail             | 21.4% |
| Little Wattlebird           | 92.9% | Grey Fantail               | 81.0% |
| Red Wattlebird              | 3.6%  | Forest Raven               | 72.6% |
| Yellow-faced Honeyeater     | 29.8% | Leaden Flycatcher          | 1.2%  |
| Spotted Pardalote           | 11.9% | Magpie-lark                | 25.0% |
| Brown Gerygone              | 92.9% | Spectacled Monarch         | 16.7% |
| Yellow-throated Scrubwren   | 48.8% | Black-faced Monarch        | 8.3%  |
| White-browed Scrubwren      | 36.9% | Rose Robin                 | 1.2%  |
| Large-billed Scrubwren      | 21.4% | Eastern Yellow Robin       | 84.5% |
| Yellow Thornbill            | 11.9% | Mistletoebird              | 10.7% |
| Striated Thornbill          | 1.2%  | Red-browed Finch           | 51.2% |
| Brown Thornbill             | 79.8% | Tree Martin                | 1.2%  |
| Black-faced Cuckoo-shrike   | 26.2% | Welcome Swallow            | 28.6% |
| White-bellied Cuckoo-shrike | 1.2%  | Silvereye                  | 56.0% |
| Golden Whistler             | 81.0% | Bassian Thrush             | 1.2%  |
| Grey Shrike-thrush          | 10.7% | Russet-tailed Thrush       | 1.2%  |
| Crested Shrike-tit          | 1.2%  |                            |       |
| Eastern Whipbird            | 83.3% |                            |       |
| Australasian Figbird        | 46.4% |                            |       |
|                             |       |                            |       |

## The 24 most commonly recorded species

| Lewin's Honeyeater      | 96.4% | Forest Raven                  | 72.6% |
|-------------------------|-------|-------------------------------|-------|
| Little Wattlebird       | 92.9% | Superb Fairy-wren             | 65.5% |
| Brown Gerygone          | 92.9% | Regent Bowerbird              | 57.1% |
| Laughing Kookaburra     | 91.7% | Silvereye                     | 56.0% |
| Eastern Yellow Robin    | 84.5% | Pied Butcherbird              | 51.2% |
| Eastern Whipbird        | 83.3% | Red-browed Finch              | 51.2% |
| Australian Magpie       | 83.3% | Silver Gull                   | 50.0% |
| Golden Whistler         | 81.0% | Australian Pied Oystercatcher | 48.8% |
| Grey Fantail            | 81.0% | Yellow-throated Scrubwren     | 48.8% |
| Australian Brush-turkey | 79.8% | Scarlet Honeyeater            | 47.6% |
| Brown Thornbill         | 79.8% | White-faced Heron             | 46.4% |
| Willie Wagtail          | 73.8% | Australasian Figbird          | 46.4% |